

# Time & Fitness–Dependent Hamiltonian Biomechanics

Tijana T. Ivancevic

Society for Nonlinear Dynamics in Human Factors, Adelaide, Australia

and

CITECH Research IP Pty Ltd, Adelaide, Australia

e-mail: tijana.ivancevic@alumni.adelaide.edu.au

## Contents

|          |                                                           |           |
|----------|-----------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                       | <b>2</b>  |
| <b>2</b> | <b>The Covariant Force Law</b>                            | <b>2</b>  |
| <b>3</b> | <b>Autonomous Hamiltonian Biomechanics</b>                | <b>5</b>  |
| <b>4</b> | <b>Time–Dependent Hamiltonian Biomechanics</b>            | <b>9</b>  |
| 4.1      | Biomechanical Bundle . . . . .                            | 9         |
| 4.2      | Biomechanical Jets . . . . .                              | 10        |
| 4.3      | Polysymplectic Dynamics . . . . .                         | 12        |
| 4.4      | Hamiltonian Connections . . . . .                         | 13        |
| 4.5      | Time–Dependent Dissipative Hamiltonian Dynamics . . . . . | 13        |
| 4.6      | Time & Fitness–Dependent Biomechanics . . . . .           | 15        |
| <b>5</b> | <b>Conclusion</b>                                         | <b>16</b> |

## Abstract

In this paper we propose the time & fitness-dependent Hamiltonian form of human biomechanics, in which *total mechanical + biochemical energy is not conserved*. Starting with the Covariant Force Law, we first develop autonomous Hamiltonian biomechanics. Then we extend it using a powerful geometrical machinery consisting of fibre bundles, jet manifolds, polysymplectic geometry and Hamiltonian connections. In this way we derive time-dependent dissipative Hamiltonian equations and the fitness evolution equation for the general time & fitness-dependent human biomechanical system.

**Keywords:** Human biomechanics, configuration bundle, Hamiltonian connections, jet manifolds, time & fitness-dependent dynamics

# 1 Introduction

Most of dynamics in both classical and quantum physics is based on *assumption of a total energy conservation* (see, e.g. [4]). Dynamics based on this assumption of time-independent energy, usually given by Hamiltonian (or Lagrangian) energy function, is called *autonomous*. This basic assumption is naturally inherited in human biomechanics, formally developed using Newton–Euler, Lagrangian or Hamiltonian formalisms (see [1, 2, 3, 5, 6, 8, 9, 10]).

And this works fine for most individual movement simulations and predictions, in which the total human energy dissipations are insignificant. However, if we analyze a 100 m-dash sprinting motion, which is in case of top athletes finished under 10 s, we can recognize a significant slow-down after about 70 m in *all* athletes – despite of their strong intention to finish and win the race, which is an obvious sign of the total energy dissipation. This can be seen, for example, in a current record-braking speeddistance curve of Usain Bolt, the world-record holder with 9.69 s, or in a former record-braking speeddistance curve of Carl Lewis, the former world-record holder (and 9 time Olympic gold medalist) with 9.86 s (see Figure 3.7 in [10]). In other words, the *total mechanical + biochemical energy* of a sprinter *cannot be conserved* even for 10 s. So, if we want to develop a realistic model of intensive human motion that is longer than 7–8 s (not to speak for instance of a 4 hour tennis match), we necessarily need to use the more advanced formalism of time-dependent mechanics.

In this paper, we will first develop the autonomous Hamiltonian biomechanics as a Hamiltonian representation of the covariant force law [see (1) in the next section] and the corresponding covariant force functor. After that we will extend the autonomous Hamiltonian biomechanics into the time-dependent one, in which *total mechanical + biochemical energy is not conserved*. for this, we will use the modern geometric formalism of jet manifolds and bundles.

## 2 The Covariant Force Law

Autonomous Hamiltonian biomechanics (as well as autonomous Lagrangian biomechanics), based on the postulate of conservation of the total mechanical energy, can be derived from the *covariant force law* [1, 2, 3, 4], which in ‘plain English’ states:

$$\text{Force 1-form} = \text{Mass distribution} \times \text{Acceleration vector-field},$$

and formally reads (using Einstein’s summation convention over repeated indices):

$$F_i = m_{ij}a^j. \tag{1}$$

Here, the force 1-form  $F_i = F_i(t, q, p) = F'_i(t, q, \dot{q})$ , ( $i = 1, \dots, n$ ) denotes any type of torques and forces acting on a human skeleton, including excitation and contraction dynamics of muscular-actuators [13, 12, 11] and rotational dynamics of hybrid robot actuators, as well as (nonlinear) dissipative joint torques and forces and external stochastic perturbation torques and forces [5].  $m_{ij}$  is the material (mass-inertia) metric tensor, which gives the total mass distribution of the human body, by including all segmental masses and their individual inertia tensors.  $a^j$  is the total acceleration vector-field, including all segmental vector-fields, defined as the absolute (Bianchi) derivative  $\dot{v}^i$  of all the segmental angular and linear velocities  $v^i = \dot{x}^i$ , ( $i = 1, \dots, n$ ), where  $n$  is the total number of active degrees of freedom (DOF) with local coordinates  $(x^i)$ .

More formally, this *central Law of biomechanics* represents the *covariant force functor*  $\mathcal{F}_*$  defined by the commutative diagram:

$$\begin{array}{ccc}
 TT^*M & \xrightarrow{\mathcal{F}_*} & TTM \\
 \uparrow F_i = \dot{p}_i & & \uparrow a^i = \dot{v}^i \\
 T^*M = \{x^i, p_i\} & & TM = \{x^i, v^i\} \\
 \swarrow p_i & & \nearrow v^i = \dot{x}^i \\
 M = \{x^i\} & & 
 \end{array} \tag{2}$$

Here,  $M \equiv M^n = \{x^i, (i = 1, \dots, n)\}$  is the biomechanical configuration  $n$ -manifold, that is the set of all active DOF of the biomechanical system under consideration (in general, human skeleton), with local coordinates  $(x^i)$ .

The right-hand branch of the fundamental covariant force functor  $\mathcal{F}_* : TT^*M \rightarrow TTM$  depicted in (2) is Lagrangian dynamics with its Riemannian geometry. To each  $n$ -dimensional ( $n$ D) smooth manifold  $M$  there is associated its  $2n$ D *velocity phase-space manifold*, denoted by  $TM$  and called the tangent bundle of  $M$ . The original configuration manifold  $M$  is called the *base* of  $TM$ . There is an onto map  $\pi : TM \rightarrow M$ , called the *projection*. Above each point  $x \in M$  there is a tangent space  $T_xM = \pi^{-1}(x)$  to  $M$  at  $x$ , which is called a fibre. The fibre  $T_xM \subset TM$  is the subset of  $TM$ , such that the total tangent bundle,  $TM = \bigsqcup_{m \in M} T_xM$ , is a disjoint union of

tangent spaces  $T_xM$  to  $M$  for all points  $x \in M$ . From dynamical perspective, the most important quantity in the tangent bundle concept is the smooth map  $v : M \rightarrow TM$ , which is an inverse to the projection  $\pi$ , i.e.,  $\pi \circ v = \text{Id}_M$ ,  $\pi(v(x)) = x$ . It is called the *velocity vector-field*  $v^i = \dot{x}^i$ .<sup>1</sup> Its graph  $(x, v(x))$  represents the cross-section of the tangent bundle  $TM$ . Velocity vector-fields are cross-sections of the tangent bundle. Biomechanical *Lagrangian* (that is, kinetic minus potential energy) is a natural energy function on the tangent bundle  $TM$ . The tangent bundle is itself a smooth manifold. It has its own tangent bundle,  $TTM$ . Cross-sections of the second tangent bundle  $TTM$  are the acceleration vector-fields.

The left-hand branch of the fundamental covariant force functor  $\mathcal{F}_* : TT^*M \rightarrow TTM$  depicted in (2) is Hamiltonian dynamics with its symplectic geometry. It takes place in the *cotangent bundle*  $T^*M_{rob}$ , defined as follows. A *dual* notion to the tangent space  $T_xM$  to a smooth manifold  $M$  at a point  $x = (x^i)$  with local is its cotangent space  $T_x^*M$  at the same point  $x$ . Similarly to the tangent bundle  $TM$ , for any smooth  $n$ D manifold  $M$ , there is associated its  $2n$ D *momentum phase-space manifold*, denoted by  $T^*M$  and called the *cotangent bundle*.  $T^*M$  is the disjoint union of all its cotangent spaces  $T_x^*M$  at all points  $x \in M$ , i.e.,  $T^*M = \bigsqcup_{x \in M} T_x^*M$ . Therefore, the cotangent

bundle of an  $n$ -manifold  $M$  is the vector bundle  $T^*M = (TM)^*$ , the (real) dual of the tangent bundle  $TM$ . Momentum 1-forms (or, covector-fields)  $p_i$  are cross-sections of the cotangent bundle. Biomechanical *Hamiltonian* (that is, kinetic plus potential energy) is a natural energy function on

<sup>1</sup>This explains the dynamical term *velocity phase-space*, given to the tangent bundle  $TM$  of the manifold  $M$ .

the cotangent bundle. The cotangent bundle  $T^*M$  is itself a smooth manifold. It has its own tangent bundle,  $TT^*M$ . Cross-sections of the mixed-second bundle  $TT^*M$  are the force 1-forms  $F_i = \dot{p}_i$ .

There is a unique smooth map from the right-hand branch to the left-hand branch of the diagram (2):

$$TM \ni (x^i, v^i) \mapsto (x^i, p^i) \in T^*M.$$

It is called the *Legendre transformation*, or *fiber derivative* (for details see, e.g. [3, 4]).

The fundamental covariant force functor  $\mathcal{F}_* : TT^*M \rightarrow TTM$  states that the force 1-form  $F_i = \dot{p}_i$ , defined on the mixed tangent-cotangent bundle  $TT^*M$ , causes the acceleration vector-field  $a^i = \dot{v}^i$ , defined on the second tangent bundle  $TTM$  of the configuration manifold  $M$ . The corresponding *contravariant acceleration functor* is defined as its inverse map,  $\mathcal{F}^* : TTM \rightarrow TT^*M$ .

Representation of human motion is rigorously defined in terms of Euclidean  $SE(3)$ -groups<sup>2</sup> of full rigid-body motion in all main human joints [6]. The configuration manifold  $M$  for human musculo-skeletal dynamics is defined as a Cartesian product of all included constrained  $SE(3)$  groups,  $M = \prod_j SE(3)^j$  where  $j$  labels the active joints. The configuration manifold  $M$  is coordinated by local joint coordinates  $x^i(t)$ ,  $i = 1, \dots, n = \text{total number of active DOF}$ . The corresponding joint velocities  $\dot{x}^i(t)$  live in the *velocity phase space*  $TM$ , which is the *tangent bundle* of the configuration manifold  $M$ .

The velocity phase-space  $TM$  has the Riemannian geometry with the *local metric form*:

$$\langle g \rangle \equiv ds^2 = g_{ij} dx^i dx^j, \quad (3)$$

where  $g_{ij}(x)$  is the material metric tensor defined by the biomechanical system's *mass-inertia matrix* and  $dx^i$  are differentials of the local joint coordinates  $x^i$  on  $M$ . Besides giving the local distances between the points on the manifold  $M$ , the Riemannian metric form  $\langle g \rangle$  defines the system's kinetic energy:

$$T = \frac{1}{2} g_{ij} \dot{x}^i \dot{x}^j,$$

giving the *Lagrangian equations* of the conservative skeleton motion with kinetic-minus-potential energy Lagrangian  $L = T - V$ , with the corresponding *geodesic form*

$$\frac{d}{dt} L_{\dot{x}^i} - L_{x^i} = 0 \quad \text{or} \quad \ddot{x}^i + \Gamma_{jk}^i \dot{x}^j \dot{x}^k = 0, \quad (4)$$

<sup>2</sup>Briefly, the Euclidean  $SE(3)$ -group is defined as a semidirect (noncommutative) product (denoted by  $\triangleright$ ) of 3D rotations and 3D translations:  $SE(3) := SO(3) \triangleright \mathbb{R}^3$ . Its most important subgroups are the following:

| Subgroup                                                                         | Definition                                                                                                                                              |
|----------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------|
| $SO(3)$ , group of rotations in 3D (a spherical joint)                           | Set of all proper orthogonal $3 \times 3$ – rotational matrices                                                                                         |
| $SE(2)$ , special Euclidean group in 2D (all planar motions)                     | Set of all $3 \times 3$ – matrices:<br>$\begin{bmatrix} \cos \theta & \sin \theta & r_x \\ -\sin \theta & \cos \theta & r_y \\ 0 & 0 & 1 \end{bmatrix}$ |
| $SO(2)$ , group of rotations in 2D subgroup of $SE(2)$ -group (a revolute joint) | Set of all proper orthogonal $2 \times 2$ – rotational matrices included in $SE(2)$ – group                                                             |
| $\mathbb{R}^3$ , group of translations in 3D (all spatial displacements)         | Euclidean 3D vector space                                                                                                                               |

where subscripts denote partial derivatives, while  $\Gamma_{jk}^i$  are the Christoffel symbols of the affine Levi-Civita connection of the biomechanical manifold  $M$ .

The corresponding momentum phase-space  $P = T^*M$  provides a natural *symplectic structure* that can be defined as follows. As the biomechanical configuration space  $M$  is a smooth  $n$ -manifold, we can pick local coordinates  $\{dx^1, \dots, dx^n\} \in M$ . Then  $\{dx^1, \dots, dx^n\}$  defines a basis of the cotangent space  $T_x^*M$ , and by writing  $\theta \in T_x^*M$  as  $\theta = p_i dx^i$ , we get local coordinates  $\{x^1, \dots, x^n, p_1, \dots, p_n\}$  on  $T^*M$ . We can now define the canonical symplectic form  $\omega$  on  $P = T^*M$  as:

$$\omega = dp_i \wedge dx^i,$$

where ‘ $\wedge$ ’ denotes the wedge or exterior product of exterior differential forms.<sup>3</sup> This 2-form  $\omega$  is obviously independent of the choice of coordinates  $\{x^1, \dots, x^n\}$  and independent of the base point  $\{x^1, \dots, x^n, p_1, \dots, p_n\} \in T_x^*M$ . Therefore, it is locally constant, and so  $d\omega = 0$ .<sup>4</sup>

If  $(P, \omega)$  is a  $2n$ D symplectic manifold then about each point  $x \in P$  there are local coordinates  $\{x^1, \dots, x^n, p_1, \dots, p_n\}$  such that  $\omega = dp_i \wedge dx^i$ . These coordinates are called canonical or symplectic. By the Darboux theorem,  $\omega$  is constant in this local chart, i.e.,  $d\omega = 0$ .

### 3 Autonomous Hamiltonian Biomechanics

We develop autonomous Hamiltonian biomechanics on the configuration biomechanical manifold  $M$  in three steps, following the standard symplectic geometry prescription (see [1, 3, 4, 7]):

**Step A** Find a symplectic *momentum phase-space*  $(P, \omega)$ .

---

<sup>3</sup>Recall that an *exterior differential form*  $\alpha$  of order  $p$  (or, a  $p$ -form  $\alpha$ ) on a base manifold  $X$  is a section of the exterior product bundle  $\overset{p}{\wedge} T^*X \rightarrow X$ . It has the following expression in local coordinates on  $X$

$$\alpha = \alpha_{\lambda_1 \dots \lambda_p} dx^{\lambda_1} \wedge \dots \wedge dx^{\lambda_p} \quad (\text{such that } |\alpha| = p),$$

where summation is performed over all ordered collections  $(\lambda_1, \dots, \lambda_p)$ .  $\Omega^p(X)$  is the vector space of  $p$ -forms on a biomechanical manifold  $X$ . In particular, the 1-forms are called the *Pfaffian forms*.

<sup>4</sup>The canonical 1-form  $\theta$  on  $T^*M$  is the unique 1-form with the property that, for any 1-form  $\beta$  which is a section of  $T^*M$  we have  $\beta^*\theta = \theta$ .

Let  $f : M \rightarrow M$  be a diffeomorphism. Then  $T^*f$  preserves the canonical 1-form  $\theta$  on  $T^*M$ , i.e.,  $(T^*f)^*\theta = \theta$ . Thus  $T^*f$  is symplectic diffeomorphism.

Recall that a symplectic structure on a smooth manifold  $M$  is a nondegenerate closed<sup>5</sup> 2-form  $\omega$  on  $M$ , i.e., for each  $x \in M$ ,  $\omega(x)$  is nondegenerate, and  $d\omega = 0$ .

Let  $T_x^*M$  be a cotangent space to  $M$  at  $m$ . The cotangent bundle  $T^*M$  represents a union  $\cup_{m \in M} T_x^*M$ , together with the standard topology on  $T^*M$  and a natural smooth manifold structure, the dimension of which is twice the dimension of  $M$ . A 1-form  $\theta$  on  $M$  represents a section  $\theta : M \rightarrow T^*M$  of the cotangent bundle  $T^*M$ .

$P = T^*M$  is our momentum phase-space. On  $P$  there is a nondegenerate symplectic 2-form  $\omega$  is defined in local joint coordinates  $x^i, p_i \in U$ ,  $U$  open in  $P$ , as  $\omega = dx^i \wedge dp_i$ . In that case the coordinates  $x^i, p_i \in U$  are called canonical. In a usual procedure the canonical 1-form  $\theta$  is first defined as  $\theta = p_i dx^i$ , and then the canonical 2-form  $\omega$  is defined as  $\omega = -d\theta$ .

A *symplectic phase-space manifold* is a pair  $(P, \omega)$ .

**Step B** Find a *Hamiltonian vector-field*  $X_H$  on  $(P, \omega)$ .

Let  $(P, \omega)$  be a symplectic manifold. A vector-field  $X : P \rightarrow TP$  is called *Hamiltonian* if there is a smooth function  $F : P \rightarrow \mathbb{R}$  such that  $i_X \omega = dF$  ( $i_X \omega$  denotes the *interior product* or *contraction* of the vector-field  $X$  and the 2-form  $\omega$ ).  $X$  is *locally Hamiltonian* if  $i_X \omega$  is closed.

Let the smooth real-valued *Hamiltonian function*  $H : P \rightarrow \mathbb{R}$ , representing the total biomechanical energy  $H(x, p) = T(p) + V(x)$  ( $T$  and  $V$  denote kinetic and potential energy of the system, respectively), be given in local canonical coordinates  $x^i, p_i \in U$ ,  $U$  open in  $P$ . The *Hamiltonian vector-field*  $X_H$ , condition by  $i_{X_H} \omega = dH$ , is actually defined via symplectic matrix  $J$ , in a local chart  $U$ , as

$$X_H = J \nabla H = (\partial_{p_i} H, -\partial_{x^i} H), \quad J = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}, \quad (5)$$

where  $I$  denotes the  $n \times n$  identity matrix and  $\nabla$  is the gradient operator.

**Step C** Find a *Hamiltonian phase-flow*  $\phi_t$  of  $X_H$ .

Let  $(P, \omega)$  be a symplectic phase-space manifold and  $X_H = J \nabla H$  a Hamiltonian vector-field corresponding to a smooth real-valued Hamiltonian function  $H : P \rightarrow \mathbb{R}$ , on it. If a unique one-parameter group of diffeomorphisms  $\phi_t : P \rightarrow P$  exists so that  $\frac{d}{dt}|_{t=0} \phi_t x = J \nabla H(x)$ , it is called the *Hamiltonian phase-flow*.

---

<sup>5</sup>A  $p$ -form  $\beta$  on a smooth manifold  $M$  is called *closed* if its exterior derivative  $d = \partial_i dx^i$  is equal to zero,

$$d\beta = 0.$$

From this condition one can see that the closed form (the *kernel* of the exterior derivative operator  $d$ ) is conserved quantity. Therefore, closed  $p$ -forms possess certain invariant properties, physically corresponding to the *conservation laws*.

Also, a  $p$ -form  $\beta$  that is an exterior derivative of some  $(p-1)$ -form  $\alpha$ ,

$$\beta = d\alpha,$$

is called *exact* (the *image* of the exterior derivative operator  $d$ ). By *Poincaré lemma*, exact forms prove to be closed automatically,

$$d\beta = d(d\alpha) = 0.$$

Since  $d^2 = 0$ , *every exact form is closed*. The converse is only partially true, by Poincaré lemma: every closed form is *locally exact*.

Technically, this means that given a closed  $p$ -form  $\alpha \in \Omega^p(U)$ , defined on an open set  $U$  of a smooth manifold  $M$  any point  $m \in U$  has a neighborhood on which there exists a  $(p-1)$ -form  $\beta \in \Omega^{p-1}(U)$  such that  $d\beta = \alpha|_U$ . In particular, there is a Poincaré lemma for contractible manifolds: Any closed form on a smoothly contractible manifold is exact.

A smooth curve  $t \mapsto (x^i(t), p_i(t))$  on  $(P, \omega)$  represents an *integral curve* of the Hamiltonian vector-field  $X_H = J\nabla H$ , if in the local canonical coordinates  $x^i, p_i \in U$ ,  $U$  open in  $P$ , *Hamiltonian canonical equations* hold:

$$\dot{q}^i = \partial_{p_i} H, \quad \dot{p}_i = -\partial_{q^i} H. \quad (6)$$

An integral curve is said to be *maximal* if it is not a restriction of an integral curve defined on a larger interval of  $\mathbb{R}$ . It follows from the standard theorem on the *existence* and *uniqueness* of the solution of a system of ODEs with smooth r.h.s, that if the manifold  $(P, \omega)$  is Hausdorff, then for any point  $x = (x^i, p_i) \in U$ ,  $U$  open in  $P$ , there exists a maximal integral curve of  $X_H = J\nabla H$ , passing for  $t = 0$ , through point  $x$ . In case  $X_H$  is complete, i.e.,  $X_H$  is  $C^p$  and  $(P, \omega)$  is compact, the maximal integral curve of  $X_H$  is the Hamiltonian phase-flow  $\phi_t : U \rightarrow U$ .

The phase-flow  $\phi_t$  is *symplectic* if  $\omega$  is constant along  $\phi_t$ , i.e.,  $\phi_t^* \omega = \omega$

( $\phi_t^* \omega$  denotes the *pull-back*<sup>6</sup> of  $\omega$  by  $\phi_t$ ),

iff  $\mathfrak{L}_{X_H} \omega = 0$

( $\mathfrak{L}_{X_H} \omega$  denotes the *Lie derivative*<sup>7</sup> of  $\omega$  upon  $X_H$ ).

Symplectic phase-flow  $\phi_t$  consists of canonical transformations on  $(P, \omega)$ , i.e., diffeomorphisms in canonical coordinates  $x^i, p_i \in U$ ,  $U$  open on all  $(P, \omega)$  which leave  $\omega$  invariant. In this case the *Liouville theorem* is valid:  $\phi_t$  *preserves* the *phase volume* on  $(P, \omega)$ . Also, the system's total energy  $H$  is conserved along  $\phi_t$ , i.e.,  $H \circ \phi_t = H$ .

Recall that the Riemannian metrics  $g = \langle, \rangle$  on the configuration manifold  $M$  is a positive-definite quadratic form  $g : TM \rightarrow \mathbb{R}$ , in local coordinates  $x^i \in U$ ,  $U$  open in  $M$ , given by (3) above. Given the metrics  $g_{ij}$ , the system's Hamiltonian function represents a momentum  $p$ -dependent quadratic form  $H : T^*M \rightarrow \mathbb{R}$  – the system's kinetic energy  $H(p) = T(p) = \frac{1}{2} \langle p, p \rangle$ , in local canonical coordinates  $x^i, p_i \in U_p$ ,  $U_p$  open in  $T^*M$ , given by

$$H(p) = \frac{1}{2} g^{ij}(x, m) p_i p_j, \quad (7)$$

where  $g^{ij}(x, m) = g_{ij}^{-1}(x, m)$  denotes the *inverse* (contravariant) material *metric tensor*

$$g^{ij}(x, m) = \sum_{\chi=1}^n m_\chi \delta_{rs} \frac{\partial x^i}{\partial x^r} \frac{\partial x^j}{\partial x^s}.$$

---

<sup>6</sup>Given a map  $f : X \rightarrow X'$  between the two manifolds, the *pullback* on  $X$  of a form  $\alpha$  on  $X'$  by  $f$  is denoted by  $f^* \alpha$ . The pullback satisfies the relations

$$f^*(\alpha \wedge \beta) = f^* \alpha \wedge f^* \beta, \quad df^* \alpha = f^*(d\alpha),$$

for any two forms  $\alpha, \beta \in \Omega^p(X)$ .

<sup>7</sup>The *Lie derivative*  $\mathfrak{L}_u \alpha$  of  $p$ -form  $\alpha$  along a vector-field  $u$  is defined by Cartan's 'magic' formula (see [3, 4]):

$$\mathfrak{L}_u \alpha = u \rfloor d\alpha + d(u \rfloor \alpha).$$

It satisfies the *Leibnitz relation*

$$\mathfrak{L}_u(\alpha \wedge \beta) = \mathfrak{L}_u \alpha \wedge \beta + \alpha \wedge \mathfrak{L}_u \beta.$$

Here, the *contraction*  $\rfloor$  of a vector-field  $u = u^\mu \partial_\mu$  and a  $p$ -form  $\alpha = \alpha_{\lambda_1 \dots \lambda_p} dx^{\lambda_1} \wedge \dots \wedge dx^{\lambda_p}$  on a biomechanical manifold  $X$  is given in local coordinates on  $X$  by

$$u \rfloor \alpha = u^\mu \alpha_{\mu \lambda_1 \dots \lambda_{p-1}} dx^{\lambda_1} \wedge \dots \wedge dx^{\lambda_{p-1}}.$$

It satisfies the following relation

$$u \rfloor (\alpha \wedge \beta) = u \rfloor \alpha \wedge \beta + (-1)^{|\alpha|} \alpha \wedge u \rfloor \beta.$$

$T^*M$  is an *orientable* manifold, admitting the standard *volume form*

$$\Omega_{\omega_H} = \frac{(-1)^{\frac{N(N+1)}{2}}}{N!} \omega_H^N.$$

For Hamiltonian vector-field,  $X_H$  on  $M$ , there is a base integral curve  $\gamma_0(t) = (x^i(t), p_i(t))$  iff  $\gamma_0(t)$  is a *geodesic*, given by the one-form *force equation*

$$\dot{\tilde{p}}_i \equiv \dot{p}_i + \Gamma_{jk}^i g^{jl} g^{km} p_l p_m = 0, \quad \text{with} \quad \dot{x}^k = g^{ki} p_i. \quad (8)$$

The l.h.s  $\dot{\tilde{p}}_i$  of the covariant momentum equation (8) represents the intrinsic or Bianchi covariant derivative of the momentum with respect to time  $t$ . Basic relation  $\dot{\tilde{p}}_i = 0$  defines the *parallel transport* on  $T^N$ , the simplest form of human-motion dynamics. In that case Hamiltonian vector-field  $X_H$  is called the *geodesic spray* and its phase-flow is called the *geodesic flow*.

For Earthly dynamics in the gravitational *potential* field  $V : M \rightarrow \mathbb{R}$ , the Hamiltonian  $H : T^*M \rightarrow \mathbb{R}$  (7) extends into potential form

$$H(p, x) = \frac{1}{2} g^{ij} p_i p_j + V(x),$$

with Hamiltonian vector-field  $X_H = J \nabla H$  still defined by canonical equations (6).

A general form of a *driven*, non-conservative Hamiltonian equations reads:

$$\dot{x}^i = \partial_{p_i} H, \quad \dot{p}_i = F_i - \partial_{x^i} H, \quad (9)$$

where  $F_i = F_i(t, x, p)$  represent any kind of joint-driving *covariant torques*, including active neuro-muscular-like controls, as functions of time, angles and momenta, as well as passive dissipative and elastic joint torques. In the covariant momentum formulation (8), the non-conservative Hamiltonian equations (9) become

$$\dot{\tilde{p}}_i \equiv \dot{p}_i + \Gamma_{jk}^i g^{jl} g^{km} p_l p_m = F_i, \quad \text{with} \quad \dot{x}^k = g^{ki} p_i.$$

The general form of autonomous Hamiltonian biomechanics is given by dissipative, driven Hamiltonian equations on  $T^*M$ :

$$\dot{x}^i = \frac{\partial H}{\partial p_i} + \frac{\partial R}{\partial p_i}, \quad (10)$$

$$\dot{p}_i = F_i - \frac{\partial H}{\partial x^i} + \frac{\partial R}{\partial x^i}, \quad (11)$$

$$x^i(0) = x_0^i, \quad p_i(0) = p_i^0, \quad (12)$$

including *contravariant equation* (10) – the *velocity vector-field*, and *covariant equation* (11) – the *force 1-form* (field), together with initial joint angles and momenta (12). Here  $R = R(x, p)$  denotes the Raileigh nonlinear (biquadratic) dissipation function, and  $F_i = F_i(t, x, p)$  are covariant driving torques of *equivalent muscular actuators*, resembling muscular excitation and contraction dynamics in rotational form. The velocity vector-field (10) and the force 1-form (11) together define the generalized Hamiltonian vector-field  $X_H$ ; the Hamiltonian energy function  $H = H(x, p)$  is its generating function.

As a Lie group, the biomechanical configuration manifold  $M = \prod_j SE(3)^j$  is Hausdorff.<sup>8</sup> Therefore, for  $x = (x^i, p_i) \in U_p$ , where  $U_p$  is an open coordinate chart in  $T^*M$ , there exists a unique one-parameter group of diffeomorphisms  $\phi_t : T^*M \rightarrow T^*M$ , that is the *autonomous Hamiltonian phase-flow*:

$$\begin{aligned} \phi_t & : T^*M \rightarrow T^*M : (p(0), x(0)) \mapsto (p(t), x(t)), \\ (\phi_t \circ \phi_s) & = \phi_{t+s}, \quad \phi_0 = \text{identity}, \end{aligned} \tag{13}$$

given by (10–12) such that

$$\left. \frac{d}{dt} \right|_{t=0} \phi_t x = J\nabla H(x).$$

## 4 Time-Dependent Hamiltonian Biomechanics

In this section we develop time-dependent Hamiltonian biomechanics. For this, we first need to extend our autonomous Hamiltonian machinery, using the general concepts of bundles, jets and connections.

### 4.1 Biomechanical Bundle

While standard autonomous Lagrangian biomechanics is developed on the configuration manifold  $X$ , the *time-dependent biomechanics* necessarily includes also the real time axis  $\mathbb{R}$ , so we have an *extended configuration manifold*  $\mathbb{R} \times X$ . Slightly more generally, the fundamental geometrical structure is the so-called *configuration bundle*  $\pi : X \rightarrow \mathbb{R}$ . Time-dependent biomechanics is thus formally developed either on the *extended configuration manifold*  $\mathbb{R} \times X$ , or on the configuration bundle  $\pi : X \rightarrow \mathbb{R}$ , using the concept of *jets*, which are based on the idea of *higher-order tangency*, or higher-order contact, at some designated point (i.e., certain anatomical joint) on a biomechanical configuration manifold  $X$ .

In general, tangent and cotangent bundles,  $TM$  and  $T^*M$ , of a smooth manifold  $M$ , are special cases of a more general geometrical object called *fibre bundle*, denoted  $\pi : Y \rightarrow X$ , where the word *fiber*  $V$  of a map  $\pi : Y \rightarrow X$  is the *preimage*  $\pi^{-1}(x)$  of an element  $x \in X$ . It is a space which *locally* looks like a product of two spaces (similarly as a manifold locally looks like Euclidean space), but may possess a different *global* structure. To get a visual intuition behind this fundamental geometrical concept, we can say that a fibre bundle  $Y$  is a *homeomorphic generalization* of a *product space*  $X \times V$  (see Figure 1), where  $X$  and  $V$  are called the *base* and the *fibre*, respectively.  $\pi : Y \rightarrow X$  is called the *projection*,  $Y_x = \pi^{-1}(x)$  denotes a fibre over a point  $x$  of the base  $X$ , while the map  $f = \pi^{-1} : X \rightarrow Y$  defines the *cross-section*, producing the *graph*  $(x, f(x))$  in the bundle  $Y$  (e.g., in case of a tangent bundle,  $f = \dot{x}$  represents a velocity vector-field).

More generally, a biomechanical configuration bundle,  $\pi : Y \rightarrow X$ , is a locally trivial fibred (or, projection) manifold over the base  $X$ . It is endowed with an atlas of fibred bundle coordinates  $(x^\lambda, y^i)$ , where  $(x^\lambda)$  are coordinates of  $X$ .

A linear connection  $\bar{\Gamma}$  on a biomechanical bundle  $Y \rightarrow X$  is given in local coordinates on  $Y$  by [14]

$$\bar{\Gamma} = dx^\lambda \otimes [\partial_\lambda - \Gamma^i_{j\lambda}(x) y^j \partial_i]. \tag{14}$$

---

<sup>8</sup>That is, for every pair of points  $x_1, x_2 \in M$ , there are disjoint open subsets (charts)  $U_1, U_2 \subset M$  such that  $x_1 \in U_1$  and  $x_2 \in U_2$ .




Figure 1: A sketch of a locally trivial fibre bundle  $Y \approx X \times V$  as a generalization of a product space  $X \times V$ ; left – main components; right – a few details (see text for explanation).

An affine connection  $\Gamma$  on a biomechanical bundle  $Y \rightarrow X$  is given in local coordinates on  $Y$  by

$$\Gamma = dx^\lambda \otimes [\partial_\lambda + (-\Gamma^i_{j\lambda}(x)y^j + \Gamma^i_\lambda(x))\partial_i].$$

Clearly, a linear connection  $\bar{\Gamma}$  is a special case of an affine connection  $\Gamma$ .

Every connection  $\Gamma$  on a biomechanical bundle  $Y \rightarrow X$  defines a system of first-order differential equations on  $Y$ , given by the local coordinate relations

$$y^i_\lambda = \Gamma^i_\lambda(y). \quad (15)$$

Integral sections for  $\Gamma$  are local solutions of (15).

## 4.2 Biomechanical Jets

A pair of smooth manifold maps,  $f_1, f_2 : M \rightarrow N$  (see Figure 2), are said to be  $k$ -tangent (or tangent of order  $k$ , or have a  $k$ th order contact) at a point  $x$  on a domain manifold  $M$ , denoted by  $f_1 \sim f_2$ , iff

$$\begin{aligned} f_1(x) &= f_2(x) && \text{called } 0\text{-tangent,} \\ \partial_x f_1(x) &= \partial_x f_2(x), && \text{called } 1\text{-tangent,} \\ \partial_{xx} f_1(x) &= \partial_{xx} f_2(x), && \text{called } 2\text{-tangent,} \\ &\dots && \text{etc. to the order } k \end{aligned}$$

In this way defined  $k$ -tangency is an *equivalence relation*.

A  $k$ -jet (or, a jet of order  $k$ ), denoted by  $j_x^k f$ , of a smooth map  $f : M \rightarrow N$  at a point  $x \in M$  (see Figure 2), is defined as an *equivalence class* of  $k$ -tangent maps at  $x$ ,

$$j_x^k f : M \rightarrow N = \{f' : f' \text{ is } k\text{-tangent to } f \text{ at } x\}.$$

For example, consider a simple function  $f : X \rightarrow Y$ ,  $x \mapsto y = f(x)$ , mapping the  $X$ -axis into the  $Y$ -axis in  $\mathbb{R}^2$ . At a chosen point  $x \in X$  we have:

a 0-jet is a graph:  $(x, f(x))$ ;

a 1-jet is a triple:  $(x, f(x), f'(x))$ ;




Figure 2: An intuitive geometrical picture behind the  $k$ -jet concept, based on the idea of a higher-order tangency (or, higher-order contact).

a 2-jet is a quadruple:  $(x, f(x), f'(x), f''(x))$ , and so on, up to the order  $k$  (where  $f'(x) = \frac{df(x)}{dx}$ , etc).

The set of all  $k$ -jets from  $j_x^k f : X \rightarrow Y$  is called the  $k$ -jet manifold  $J^k(X, Y)$ .

Formally, given a biomechanical bundle  $Y \rightarrow X$ , its first-order *jet manifold*  $J^1 Y$  comprises the set of equivalence classes  $j_x^1 s$ ,  $x \in X$ , of sections  $s : X \rightarrow Y$  so that sections  $s$  and  $s'$  belong to the same class iff

$$Ts|_{T_x X} = Ts'|_{T_x X}.$$

Intuitively, sections  $s, s' \in j_x^1 s$  are identified by their values  $s^i(x) = s'^i(x)$  and the values of their partial derivatives  $\partial_\mu s^i(x) = \partial_\mu s'^i(x)$  at the point  $x$  of  $X$ . There are the natural fibrations [14]

$$\pi_1 : J^1 Y \ni j_x^1 s \mapsto x \in X, \quad \pi_{01} : J^1 Y \ni j_x^1 s \mapsto s(x) \in Y.$$

Given bundle coordinates  $(x^\lambda, y^i)$  of  $Y$ , the associated jet manifold  $J^1 Y$  is endowed with the adapted coordinates

$$(x^\lambda, y^i, y_\lambda^i), \quad (y^i, y_\lambda^i)(j_x^1 s) = (s^i(x), \partial_\lambda s^i(x)), \quad y_\lambda^i = \frac{\partial x^\mu}{\partial x'^\lambda} (\partial_\mu + y_\mu^j \partial_j) y^i.$$

In particular, given the biomechanical configuration bundle  $M \rightarrow \mathbb{R}$  over the time axis  $\mathbb{R}$ , the 1-jet space  $J^1(\mathbb{R}, M)$  is the set of equivalence classes  $j_t^1 s$  of sections  $s^i : \mathbb{R} \rightarrow M$  of the configuration bundle  $M \rightarrow \mathbb{R}$ , which are identified by their values  $s^i(t)$ , as well as by the values of their partial derivatives  $\partial_t s^i = \partial_t s^i(t)$  at time points  $t \in \mathbb{R}$ . The 1-jet manifold  $J^1(\mathbb{R}, M)$  is coordinated by  $(t, x^i, \dot{x}^i)$ , that is by *(time, coordinates and velocities)* at every active human joint, so the 1-jets are local joint coordinate maps

$$j_t^1 s : \mathbb{R} \rightarrow M, \quad t \mapsto (t, x^i, \dot{x}^i).$$

The *repeated jet manifold*  $J^1 J^1 Y$  is defined to be the jet manifold of the bundle  $J^1 Y \rightarrow X$ . It is endowed with the adapted coordinates  $(x^\lambda, y^i, y_\lambda^i, y_{\lambda\mu}^i, y_{\lambda\mu}^i)$ .

The *second-order jet manifold*  $J^2 Y$  of a bundle  $Y \rightarrow X$  is the subbundle of  $\widehat{J}^2 Y \rightarrow J^1 Y$  defined by the coordinate conditions  $y_{\lambda\mu}^i = y_{\mu\lambda}^i$ . It has the local coordinates  $(x^\lambda, y^i, y_\lambda^i, y_{\lambda\leq\mu}^i)$

together with the transition functions [14]

$$y'^i_{\lambda\mu} = \frac{\partial x^\alpha}{\partial x'^\mu} (\partial_\alpha + y^j_\alpha \partial_j + y^\nu_{\nu\alpha} \partial^\nu_j) y'^i_\lambda.$$

The second-order jet manifold  $J^2Y$  of  $Y$  comprises the equivalence classes  $j^2_x s$  of sections  $s$  of  $Y \rightarrow X$  such that

$$y^i_\lambda(j^2_x s) = \partial_\lambda s^i(x), \quad y^i_{\lambda\mu}(j^2_x s) = \partial_\mu \partial_\lambda s^i(x).$$

In other words, two sections  $s, s' \in j^2_x s$  are identified by their values and the values of their first and second-order derivatives at the point  $x \in X$ .

In particular, given the biomechanical configuration bundle  $M \rightarrow \mathbb{R}$  over the time axis  $\mathbb{R}$ , the 2-jet space  $J^2(\mathbb{R}, M)$  is the set of equivalence classes  $j^2_t s$  of sections  $s^i : \mathbb{R} \rightarrow M$  of the configuration bundle  $\pi : M \rightarrow \mathbb{R}$ , which are identified by their values  $s^i(t)$ , as well as the values of their first and second partial derivatives,  $\partial_t s^i = \partial_t s^i(t)$  and  $\partial_{tt} s^i = \partial_{tt} s^i(t)$ , respectively, at time points  $t \in \mathbb{R}$ . The 2-jet manifold  $J^2(\mathbb{R}, M)$  is coordinated by  $(t, x^i, \dot{x}^i, \ddot{x}^i)$ , that is by (*time, coordinates, velocities and accelerations*) at every active human joint, so the 2-jets are local joint coordinate maps<sup>9</sup>

$$j^2_t s : \mathbb{R} \rightarrow M, \quad t \mapsto (t, x^i, \dot{x}^i, \ddot{x}^i).$$

### 4.3 Polysymplectic Dynamics

Let  $Y \rightarrow X$  be a biomechanical bundle with local coordinates  $(x^\lambda, y^i)$ . In jet terms, a *first-order Lagrangian* is defined (through the standard Lagrangian density  $\mathcal{L}$ ) to be a horizontal density  $L = \mathcal{L}\omega$  on the jet manifold  $J^1Y$ . The jet manifold  $J^1Y$  plays the role of the finite-dimensional configuration space of sections of the bundle  $Y \rightarrow X$ .

Lagrangian  $L$  yields the *Legendre morphism*  $\widehat{L} : J^1Y \rightarrow \Pi$ , from the 1-jet manifold  $J^1Y$  to the *Legendre manifold*, given by a product [14]

$$\Pi = V^*Y \wedge (\wedge^{n-1} T^*X) = V^*Y \wedge (\wedge^n T^*X) \otimes TX, \quad (16)$$

where  $V$  is called the *vertical bundle*.  $\Pi$  plays the role of the finite-dimensional phase-space of sections of  $Y \rightarrow X$ . Given the bundle coordinates  $(x^\lambda, y^i)$  on  $Y \rightarrow X$ , the Legendre bundle (16) has local coordinates  $(x^\lambda, y^i, p^\lambda_i)$ , where  $p^\lambda_i$  are the holonomic coordinates with the transition functions

$$p'^\lambda_i = \det\left(\frac{\partial x^\epsilon}{\partial x'^\nu}\right) \frac{\partial y^j}{\partial y'^i} \frac{\partial x'^\lambda}{\partial x^\mu} p^\mu_j. \quad (17)$$

Relative to these coordinates, the Legendre morphism  $\widehat{L}$  reads

$$p^\mu_i \circ \widehat{L} = \pi^\mu_i. \quad (18)$$

The Legendre manifold  $\Pi$  in (16) is equipped with the *generalized Liouville form*

$$\Theta = -p^\lambda_i dy^i \wedge \omega \otimes \partial_\lambda, \quad (19)$$

---

<sup>9</sup>For more technical details on jet spaces with their physical applications, see [14, 15]).

where  $\otimes$  denotes the standard tensor product. Furthermore, for any Pfaffian form  $\theta$  on  $X$  we have the relation

$$\Lambda \rfloor \theta = -d(\Theta \rfloor \theta).$$

This relation introduces the the *canonical polysymplectic form* on the Legendre manifold  $\Pi$ ,

$$\Lambda = dp_i^\lambda \wedge dy^i \wedge \omega \otimes \partial_\lambda, \quad (20)$$

whose coordinate expression (20) is maintained under holonomic coordinate transformations (17). It is a pullback-valued form [14].

#### 4.4 Hamiltonian Connections

Let  $J^1\Pi$  be the jet manifold of the *Legendre bundle*  $\Pi \rightarrow X$ . It is endowed with the adapted coordinates  $(x^\lambda, y^i, p_i^\lambda, y_\mu^i, p_{i\mu}^\lambda)$ . By analogy with the notion of an autonomous Hamiltonian vector-field  $X_H$  in (5), a connection  $\gamma$  on the bundle  $\Pi \rightarrow X$ , given by

$$\gamma = dx^\lambda \otimes (\partial_\lambda + \gamma_\lambda^i \partial_i + \gamma_{i\lambda}^\mu \partial_\mu^i),$$

is said to be *locally Hamiltonian* if the exterior form  $\gamma \rfloor \Lambda$  is closed and *Hamiltonian* if the form  $\gamma \rfloor \Lambda$  is exact. A connection  $\gamma$  is locally Hamiltonian iff it obeys the conditions [14, 15]

$$\partial_\lambda^i \gamma_\mu^j - \partial_\mu^j \gamma_\lambda^i = 0, \quad \partial_i \gamma_{j\mu}^\mu - \partial_j \gamma_{i\mu}^\mu = 0, \quad \partial_j \gamma_\lambda^i + \partial_\lambda^i \gamma_{j\mu}^\mu = 0. \quad (21)$$

A  $p$ -form  $H$  on the Legendre bundle  $\Pi \rightarrow X$  is called a *general Hamiltonian form* if there exists a Hamiltonian connection such that

$$\gamma \rfloor \Lambda = dH.$$

A general, dissipative, time-dependent Hamiltonian form  $H$  on  $\Pi$  is said to be *Hamiltonian* if it has the splitting

$$H = H_\Gamma - \tilde{H}_\Gamma = p_i^\lambda dy^i \wedge \omega_\lambda - (p_i^\lambda \Gamma_\lambda^i + \tilde{\mathcal{H}}_\Gamma) \omega = p_i^\lambda dy^i \wedge \omega_\lambda - \mathcal{H} \omega \quad (22)$$

modulo closed forms, where  $\Gamma$  is a connection on  $Y$  and  $\tilde{H}_\Gamma$  is a horizontal density. This splitting is preserved under the holonomic coordinate transformations (17).

Let a Hamiltonian connection  $\gamma$  associated with a Hamiltonian form  $H$  have an integral section  $s$  of  $\Pi \rightarrow X$ , that is,  $\gamma \circ s = J^1 s$ . Then  $s$  satisfies the system of first-order *Hamiltonian equations* on  $\Pi$ ,

$$y_\lambda^i = \partial_\lambda^i \mathcal{H}, \quad p_{i\lambda}^\lambda = -\partial_i \mathcal{H}.$$

#### 4.5 Time-Dependent Dissipative Hamiltonian Dynamics

We can now formulate the time-dependent biomechanics as an  $n = 1$ -reduction of polysymplectic dynamics described above. The biomechanical phase space is the Legendre manifold  $\Pi$ , endowed with the holonomic coordinates  $(t, y^i, p_i)$  with the transition functions

$$p_i' = \frac{\partial y^j}{\partial y'^i} p_j.$$

$\Pi$  admits the canonical form  $\Lambda$  (20), which now reads

$$\Lambda = dp_i \wedge dy^i \wedge dt \otimes \partial_t.$$

As a particular case of the polysymplectic machinery of the previous section, we say that a connection

$$\gamma = dt \otimes (\partial_t + \gamma^i \partial_i + \gamma_i \partial^i)$$

on the bundle  $\Pi \rightarrow X$  is locally Hamiltonian if the exterior form  $\gamma \rfloor \Lambda$  is closed and Hamiltonian if the form  $\gamma \rfloor \Lambda$  is exact. A connection  $\gamma$  is locally Hamiltonian iff it obeys the conditions (21) which now take the form

$$\partial^i \gamma^j - \partial^j \gamma^i = 0, \quad \partial_i \gamma_j - \partial_j \gamma_i = 0, \quad \partial_j \gamma^i + \partial^i \gamma_j = 0.$$

Note that every connection  $\Gamma = dt \otimes (\partial_t + \Gamma^i \partial_i)$  on the bundle  $Y \rightarrow X$  gives rise to the Hamiltonian connection  $\tilde{\Gamma}$  on  $\Pi \rightarrow X$ , given by

$$\tilde{\Gamma} = dt \otimes (\partial_t + \Gamma^i \partial_i - \partial_j \Gamma^i p_i \partial^j).$$

The corresponding Hamiltonian form  $H_\Gamma$  is given by

$$H_\Gamma = p_i dy^i - p_i \Gamma^i dt.$$

Let  $H$  be a dissipative Hamiltonian form (22) on  $\Pi$ , which reads:

$$H = p_i dy^i - \mathcal{H} dt = p_i dy^i - p_i \Gamma^i dt - \tilde{\mathcal{H}}_\Gamma dt. \quad (23)$$

We call  $\mathcal{H}$  and  $\tilde{\mathcal{H}}$  in the decomposition (23) the *Hamiltonian* and the *Hamiltonian function* respectively. Let  $\gamma$  be a Hamiltonian connection on  $\Pi \rightarrow X$  associated with the Hamiltonian form (23). It satisfies the relations [14, 15]

$$\begin{aligned} \gamma \rfloor \Lambda &= dp_i \wedge dy^i + \gamma_i dy^i \wedge dt - \gamma^i dp_i \wedge dt = dH, \\ \gamma^i &= \partial^i \mathcal{H}, \quad \gamma_i = -\partial_i \mathcal{H}. \end{aligned} \quad (24)$$

From equations (24) we see that, in the case of biomechanics, one and only one Hamiltonian connection is associated with a given Hamiltonian form.

Every connection  $\gamma$  on  $\Pi \rightarrow X$  yields the system of first-order differential equations (15), which now takes the explicit form:

$$\dot{y}^i = \gamma^i, \quad \dot{p}_i = \gamma_i. \quad (25)$$

They are called the *evolution equations*. If  $\gamma$  is a Hamiltonian connection associated with the Hamiltonian form  $H$  (23), the evolution equations (25) become the *dissipative time-dependent Hamiltonian equations*:

$$\dot{y}^i = \partial^i \mathcal{H}, \quad \dot{p}_i = -\partial_i \mathcal{H}. \quad (26)$$

In addition, given any scalar function  $f$  on  $\Pi$ , we have the *dissipative Hamiltonian evolution equation*

$$d_H f = (\partial_t + \partial^i \mathcal{H} \partial_i - \partial_i \mathcal{H} \partial^i) f, \quad (27)$$

relative to the Hamiltonian  $\mathcal{H}$ . On solutions  $s$  of the Hamiltonian equations (26), the evolution equation (27) is equal to the total time derivative of the function  $f$ :

$$s^* d_H f = \frac{d}{dt} (f \circ s).$$

## 4.6 Time & Fitness-Dependent Biomechanics

The dissipative Hamiltonian system (26)–(27) is the basis for our time & fitness-dependent biomechanics. The scalar function  $f$  in (27) on the biomechanical Legendre phase-space manifold  $\Pi$  is now interpreted as an *individual neuro-muscular fitness function*. This fitness function is a ‘determinant’ for the performance of muscular drives for the driven, dissipative Hamiltonian biomechanics. These muscular drives, for all active DOF, are given by time & fitness-dependent Pfaffian form:  $F_i = F_i(t, y, p, f)$ . In this way, we obtain our final model for time & fitness-dependent Hamiltonian biomechanics:

$$\begin{aligned}\dot{y}^i &= \partial^i \mathcal{H}, \\ \dot{p}_i &= F_i - \partial_i \mathcal{H}, \\ d_H f &= (\partial_t + \partial^i \mathcal{H} \partial_i - \partial_i \mathcal{H} \partial^i) f.\end{aligned}$$

Physiologically, the active muscular drives  $F_i = F_i(t, y, p, f)$  consist of [1, 2]):

**1. Synovial joint mechanics**, giving the first stabilizing effect to the conservative skeleton dynamics, is described by the  $(y, \dot{y})$ -form of the *Rayleigh–Van der Pol’s dissipation function*

$$R = \frac{1}{2} \sum_{i=1}^n (\dot{y}^i)^2 [\alpha_i + \beta_i (y^i)^2],$$

where  $\alpha_i$  and  $\beta_i$  denote dissipation parameters. Its partial derivatives give rise to the viscous-damping torques and forces in the joints

$$\mathcal{F}_i^{joint} = \partial R / \partial \dot{y}^i,$$

which are linear in  $\dot{y}^i$  and quadratic in  $y^i$ .

**2. Muscular mechanics**, giving the driving torques and forces  $\mathcal{F}_i^{musc} = \mathcal{F}_i^{musc}(t, y, \dot{y})$  with  $(i = 1, \dots, n)$  for human biomechanics, describes the internal excitation and contraction dynamics of *equivalent muscular actuators* [11].

(a) The *excitation dynamics* can be described by an impulse force–time relation

$$\begin{aligned}F_i^{imp} &= F_i^0 (1 - e^{-t/\tau_i}) & \text{if stimulation} > 0 \\ F_i^{imp} &= F_i^0 e^{-t/\tau_i} & \text{if stimulation} = 0,\end{aligned}$$

where  $F_i^0$  denote the maximal isometric muscular torques and forces, while  $\tau_i$  denote the associated time characteristics of particular muscular actuators. This relation represents a solution of the Wilkie’s muscular *active-state element* equation [12]

$$\dot{\mu} + \Gamma \mu = \Gamma S A, \quad \mu(0) = 0, \quad 0 < S < 1,$$

where  $\mu = \mu(t)$  represents the active state of the muscle,  $\Gamma$  denotes the element gain,  $A$  corresponds to the maximum tension the element can develop, and  $S = S(r)$  is the ‘desired’ active state as a

function of the motor unit stimulus rate  $r$ . This is the basis for biomechanical force controller.

(b) The *contraction dynamics* has classically been described by the Hill's *hyperbolic force-velocity* relation [13]

$$F_i^{Hill} = \frac{(F_i^0 b_i - \delta_{ij} a_i \dot{y}^j)}{(\delta_{ij} \dot{y}^j + b_i)},$$

where  $a_i$  and  $b_i$  denote the Hill's parameters, corresponding to the energy dissipated during the contraction and the phosphagenic energy conversion rate, respectively, while  $\delta_{ij}$  is the Kronecker's  $\delta$ -tensor.

In this way, human biomechanics describes the excitation/contraction dynamics for the  $i$ th equivalent muscle-joint actuator, using the simple impulse-hyperbolic product relation

$$\mathcal{F}_i^{musc}(t, y, \dot{y}) = F_i^{imp} \times F_i^{Hill}.$$

Now, for the purpose of biomedical engineering and rehabilitation, human biomechanics has developed the so-called *hybrid rotational actuator*. It includes, along with muscular and viscous forces, the D.C. motor drives, as used in robotics

$$\begin{aligned} \mathcal{F}_k^{robo} &= i_k(t) - J_k \ddot{y}_k(t) - B_k \dot{y}_k(t), \quad \text{with} \\ l_k i_k(t) + R_k i_k(t) + C_k \dot{y}_k(t) &= u_k(t), \end{aligned}$$

where  $k = 1, \dots, n$ ,  $i_k(t)$  and  $u_k(t)$  denote currents and voltages in the rotors of the drives,  $R_k, l_k$  and  $C_k$  are resistances, inductances and capacitances in the rotors, respectively, while  $J_k$  and  $B_k$  correspond to inertia moments and viscous dampings of the drives, respectively.

Finally, to make the model more realistic, we need to add some *stochastic torques and forces*:

$$\mathcal{F}_i^{stoch} = B_{ij}[y^i(t), t] dW^j(t),$$

where  $B_{ij}[y(t), t]$  represents continuous stochastic *diffusion fluctuations*, and  $W^j(t)$  is an  $N$ -variable *Wiener process* (i.e., generalized Brownian motion) [5], with

$$dW^j(t) = W^j(t + dt) - W^j(t), \quad (\text{for } j = 1, \dots, n = \text{no. of active DOF}).$$

## 5 Conclusion

We have presented the time-dependent generalization of an 'ordinary' autonomous Hamiltonian biomechanics, in which total mechanical + biochemical energy is not conserved. Starting with the Covariant Force Law, we have first developed autonomous Hamiltonian biomechanics. Then we have introduced a general framework for time-dependent Hamiltonian biomechanics in terms jets, Legendre manifolds and dissipative Hamiltonian connections associated with the extended musculo-skeletal configuration manifold, called the configuration bundle. In this way we formulated a general Hamiltonian model for time & fitness-dependent human biomechanics.

## References

- [1] Ivancevic, V., Ivancevic, T., Human–Like Biomechanics: A Unified Mathematical Approach to Human Biomechanics and Humanoid Robotics. Springer, Dordrecht, (2006)
- [2] Ivancevic, V., Ivancevic, T., Natural Biodynamics. World Scientific, Singapore (2006)
- [3] Ivancevic, V., Ivancevic, T., Geometrical Dynamics of Complex Systems: A Unified Modelling Approach to Physics, Control, Biomechanics, Neurodynamics and Psycho-Socio-Economical Dynamics. Springer, Dordrecht, (2006)
- [4] Ivancevic, V., Ivancevic, T., Applied Differential Geometry: A Modern Introduction. World Scientific, Singapore, (2007)
- [5] Ivancevic, V., Ivancevic, T., High–Dimensional Chaotic and Attractor Systems. Springer, Berlin, (2006)
- [6] Ivancevic, V., Ivancevic, T., Human versus humanoid robot biodynamcis. *Int. J. Hum. Rob.* **5**(4), 699–713, (2008)
- [7] Ivancevic, V., Ivancevic, T., Complex Nonlinearity: Chaos, Phase Transitions, Topology Change and Path Integrals, Springer, Berlin, (008)
- [8] Ivancevic, T., Jovanovic, B., Djukic, M., Markovic, S., Djukic, N., Biomechanical Analysis of Shots and Ball Motion in Tennis and the Analogy with Handball Throws, *J. Facta Universitatis, Series: Sport*, **6**(1), 51–66, (2008)
- [9] Ivancevic, T., Jain, L., Pattison, J., Hariz, A., Nonlinear Dynamics and Chaos Methods in Neurodynamics and Complex Data Analysis, *Nonl. Dyn. (Springer)*, **56**(1-2), 23–44, (2009)
- [10] Ivancevic, T., Jovanovic, B., Djukic, S., Djukic, M., Markovic, S., Complex Sports Biodynamics: With Practical Applications in Tennis, Springer, Berlin, (2009)
- [11] Hatze, H., A general myocybernetic control model of skeletal muscle. *Biol. Cyber.* **28**, 143–157, (1978)
- [12] Wilkie, D.R., The mechanical properties of muscle. *Brit. Med. Bull.* **12**, 177–182, (1956)
- [13] Hill, A.V., The heat of shortening and the dynamic constants of muscle. *Proc. Roy. Soc.* **B76**, 136–195, (1938)
- [14] Giachetta, G., Mangiarotti, L., Sardanashvily, G., New Lagrangian and Hamiltonian Methods in Field Theory, World Scientific, Singapore, (1997)
- [15] Sardanashvily, G.: Hamiltonian time-dependent mechanics. *J. Math. Phys.* **39**, 2714, (1998)